



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

On entropy scaling laws for diffusion

Sorin Bastea

June 7, 2004

Physical Review Letters

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

On entropy scaling laws for diffusion

Sorin Bastea*

Lawrence Livermore National Laboratory,

P.O. BOX 808, Livermore, CA 94550

PACS numbers: 66.10.Cb

In a recent letter [1], Samanta, Musharaf Ali and Ghosh set out to understand the low density failure of the entropy scaling law for the self-diffusion coefficient D conjectured by Dzugutov [2] and to provide a simple alternative. After an interesting derivation, that however contains a number of uncontrolled approximations, they arrive at Eq. 7 of [1], which reduces for a hard sphere fluid to:

$$\frac{D}{D_E} = \frac{A}{1 - s_e/k_B} \quad (1)$$

where $D_E = D_B/\chi$ is the Enskog diffusion coefficient [3], $D_B = 3(k_B T/\pi m)^{1/2}/8\rho\sigma^2$ the Boltzmann diffusion coefficient, χ the contact value of the pair correlation function, s_e the excess entropy per particle, and A a constant that the authors set to 2.5. Using the excess entropy $s_e/k_B = -(4\eta - 3\eta^2)/(1 - \eta)^2$ and contact value $\chi = (2 - \eta)/2(1 - \eta)^3$ obtained from the Carnahan-Starling equation of state [4] ($\eta = \pi\rho\sigma^3/6$ - packing fraction), we test this relation against the hard sphere molecular dynamics simulation results of Erpenbeck and Wood (EW)[5] - Fig. 1. The disagreement is quite significant and more important the behavior of the two curves is very different, as $(D/D_E)_{EW}$ is not a monotonically decreasing function of $(-s_e/k_B)$. The discrepancy cannot be attributed to the authors use of an approximation for the excess entropy.

The idea of entropy scaling for transport coefficients has a fairly long history [6]. Arguing on the basis of the molecular “caging” effect Dzugutov [2] proposed the scaling law:

$$\frac{D}{\sigma^2\Gamma_E} = B \exp(s_e/k_B) \quad (2)$$

which assumes that the natural length and time scales for diffusion are given by a suitably defined hard sphere diameter σ and the Enskog collision frequency $\Gamma_E = 4\sigma^2\chi\rho\sqrt{\pi k_B T/m}$, with B an universal constant. Unfortunately, the above relation appears to work only in a limited, high density domain for both hard spheres [7] and realistically modeled fluids [8]. The problem that arises at low and moderate densities (see Fig. 1) with the scaling introduced in Eq. 2 can be understood if we observe that the left-hand-side of that equation can be written up to a multiplicative constant as $D/D_B\chi\eta^2$. Therefore, in the limit of a dilute system, $\eta \rightarrow 0$, this term diverges as $1/\eta^2$, while the right-hand-side of Eq. 2 remains finite. This behavior should be expected for any valid definition of σ and χ and s_e approximation.

The noted pathology of Eq. 2 can be avoided by replacing σ as the relevant length scale with $1/\rho\sigma^2$, the Boltzmann mean-free path, which should be a reasonable measure of the

degree of molecular confinement, $1/\rho\sigma^2 \propto \sigma/\eta$. The new relationship is:

$$\frac{D}{D_{B\chi}} = \exp(\gamma s_e/k_B) \quad (3)$$

where we introduced a different constant γ . The test of this suggested dependence is shown in Fig. 2 for hard spheres, with $\gamma = 0.8$. Furthermore, Eq. 3 holds for Van der Waals fluids as well when a reasonable definition for σ is used [8].

Samanta et al. also propose a generalized Stokes-Einstein relation connecting the diffusion coefficient and the viscosity. It is worth pointing out that such a relation is hardly necessary given that the usual Stokes-Einstein relation with the 'slip' boundary condition holds well for both hard spheres [9] and Van der Waals fluids [8].

This work was performed under the auspices of the U. S. Department of Energy by University of California Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48.

* Electronic address: `bastea2@llnl.gov`

- [1] A. Samanta, Sk. Musharaf Ali, S.K. Ghosh, Phys. Rev. Lett. **92**, 145901 (2004).
- [2] M. Dzugutov, Nature(London) **381**, 137 (1996).
- [3] S. Chapman, T.G. Cowling, *The Mathematical Theory of Non-uniform Gases* (Cambridge University Press, Cambridge, England, 1970).
- [4] J.-P. Hansen, I.R. McDonald, *Theory of Simple Liquids*, 2nd edition, (Academic Press, London, 1986).
- [5] J.J. Erpenbeck, W.W. Wood, Phys. Rev. A **43**, 4254 (1991).
- [6] Y. Rosenfeld, Phys. Rev. A **15**, 2545 (1977).
- [7] E.G.D. Cohen, L. Rondoni, Phys. Rev. Lett. **84**, 394 (2000).
- [8] S. Bastea, Phys. Rev. E **68**, 031204 (2003).
- [9] B.J. Alder, D.M. Gass, T.E. Wainwright, J. Chem. Phys. **53**, 3813 (1970).




FIG. 1: Comparison of the hard spheres diffusion coefficient (simulation results of Ref. [5]) - circles, with the scaling relation proposed by Samanta et al. [1] (Eq. 1) - dashed line, Dzugutov scaling law [2] (Eq. 2) - dot-dashed line, and new entropy scaling (Eq. 3) - solid line.